

## Product Datasheet

# Rabbit IgG (H&L) Antibody Biotin Conjugated Pre-Adsorbed (orb347727)

<b>Catalog Number</b>	orb347727
<b>Category</b>	Antibodies
<b>Description</b>	Rabbit IgG (H&L) antibody (Biotin)
<b>Clonality</b>	Polyclonal
<b>Species/Host</b>	Mouse
<b>Isotype</b>	IgG
<b>Conjugation</b>	Biotin
<b>Reactivity</b>	Rabbit
<b>Form/Appearance</b>	Lyophilized
<b>Concentration</b>	1.0 mg/mL
<b>Buffer/Preservatives</b>	Preservative: 0.01% (w/v) Sodium Azide. Stabilizer: 10 mg/mL Bovine Serum Albumin (rAlbumin) - Immunoglobulin and Protease free; Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
<b>Purity</b>	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Mouse Serum, Rabbit IgG and Rabbit Serum. No reaction was observed against Human, Goat and Mouse Serum Proteins.
<b>Immunogen</b>	Rabbit IgG whole molecule

**Biorbyt Ltd.**

7 Signet Court, Swann Road  
Cambridge  
CB5 8LA  
United Kingdom


Email: [info@biorbyt.com](mailto:info@biorbyt.com), [support@biorbyt.com](mailto:support@biorbyt.com)  
Phone: [+44 \(0\)1223 859353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)

**Biorbyt LLC**

68 TW Alexander Drive  
Research Triangle Park  
Durham  
NC 27713  
United States

Email: [info@biorbyt.com](mailto:info@biorbyt.com), [support@biorbyt.com](mailto:support@biorbyt.com)  
Phone: [+1 \(415\) 906-5211](tel:+1(415)906-5211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)

<b>Tested applications</b>	ELISA, IHC, WB
<b>Dilution range</b>	ELISA: 1:20,000 - 1:100,000, IHC: 1:1,000 - 1:5,000, WB: 1:2,000 - 1:10,000
<b>Application notes</b>	Mouse Anti-Rabbit IgG Biotin Conjugate has been assayed against 1.0 ug of Rabbit IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin and ABTS (2,2'-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid]) code as a substrate for 30 minutes at room temperature. A working dilution of 1:14,000 to 1:60,000 is suggested for this product. Reconstitution Buffer: Restore with deionized water (or equivalent). Reconstitution Volume: 1.0 mL
<b>Antibody Type</b>	Secondary Antibody
<b>Storage</b>	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
<b>Note</b>	For research use only
<b>Expiration Date</b>	12 months from date of receipt.



GRK5 regulates dendritic development. (A and B) Hippocampal neuron cultures transfected at DIV5 were observed at DIV8. Total dendritic branch tip numbers (TDBTN) and total dendrite length of transfected neurons were measured. For each group, 40–60 (A) or 30–40 (B) neurons from three independent cultures were analyzed. One-way ANOVA followed by Tukey–Kramer posthoc test. (C and D) Hippocampal neurons were transfected at DIV9 and observed at DIV17. Boxed regions are enlarged below each image. For each group, 30–40 dendrites of 8–10 neurons from three independent cultures were analyzed. Protrusion and spine number were measured. (C) GFP was cotransfected with GRK5 variants to visualize dendritic spines (one-way ANOVA followed by Tukey–Kramer posthoc test). (D) Neuron cultures transfected with control or GRK5 RNAi constructs (Student's t test). Bars, 10  $\mu$ m. Error bars indicate SEM. \*, P 0.03; \*\*, P 0.01; \*\*\*, P 0.001. Ctrl, control.

**Biorbyt Ltd.**

7 Signet Court, Swann Road  
Cambridge  
CB5 8LA  
United Kingdom

Email: [info@biorbyt.com](mailto:info@biorbyt.com), [support@biorbyt.com](mailto:support@biorbyt.com)

Phone: +44 (0)1223 859353 | Fax: +1 (415) 651-8558


**Biorbyt LLC**

68 TW Alexander Drive  
Research Triangle Park  
Durham  
NC 27713  
United States

Email: [info@biorbyt.com](mailto:info@biorbyt.com), [support@biorbyt.com](mailto:support@biorbyt.com)

Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558

Mouse Anti-Rabbit IgG biotin conjugated antibody. Peri-Urethral Prostate Tissues Exhibit Fibroblastic and Myofibroblastic Cell Populations. Peri-urethral prostate tissues from patients 1007 and 0516 were explanted and primary fibroblasts were isolated and grown to monolayer cultures. Photomicrographs demonstrate fibroblastic morphology for 0516 primary cells but mixed fibroblastic and myofibroblastic morphologies for patient 1007. Cells from both cultures were then stained for collagen 1 (COL1) (PE-cy5-conjugated Ab, red),  $\alpha$ -smooth muscle actin ( $\alpha$ SMA) (fluorescein-conjugated Ab, green), or the nuclei counterstained with DAPI (blue). Merged images show that primary cells from patient 1007 exhibited high levels of co-localized COL1 and  $\alpha$ SMA protein expression (yellow) consistent with a myofibroblastic phenotype. Control mouse IgG2a and rabbit IgG biotin conjugate were used at 1:2000 dilution. All images were captured at 400X in visible light on brightfield settings.

**Biorbyt Ltd.**

7 Signet Court, Swann Road  
Cambridge  
CB5 8LA  
United Kingdom

Email: [info@biorbyt.com](mailto:info@biorbyt.com), [support@biorbyt.com](mailto:support@biorbyt.com)

Phone: [+44 \(0\)1223 859353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)

**Biorbyt LLC**

68 TW Alexander Drive  
Research Triangle Park  
Durham  
NC 27713  
United States

Email: [info@biorbyt.com](mailto:info@biorbyt.com), [support@biorbyt.com](mailto:support@biorbyt.com)

Phone: [+1 \(415\) 906-5211](tel:+1(415)906-5211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)