

## **Product Datasheet**

# Sheep IgG Fc Antibody Peroxidase Conjugated (orb347925)

Catalog Number orb347925

**Category** Antibodies

**Description** Sheep IgG F(c) antibody (Peroxidase)

**Clonality** Polyclonal

Species/Host Rabbit

**Isotype** lgG

**Conjugation** HRP

**Reactivity** Sheep

Form/Appearance Lyophilized

**Concentration** 1.5 mg/mL

**Buffer/Preservatives** 0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!

**Purity** This product was prepared from monospecific antiserum by immunoaffinity

chromatography using Sheep IgG coupled to agarose. Assay by

immunoelectrophoresis resulted in a single precipitin arc against anti-

Peroxidase, anti-Rabbit Serum, Sheep IgG, Sheep IgG F(c) and Sheep Serum. No

reaction was observed against Sheep IgG F(ab).

**Immunogen** Anti-Sheep IgG was produced by repeated immunization with sheep IgG F(c)

fragment in rabbit.

**Tested applications** ELISA, IHC, WB

### **Biorbyt Ltd.**

7 Signet Court, Swann Road Cambridge CB5 8LA United Kingdom

Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: <u>+44 (0)1223 859353</u> | Fax: <u>+1 (415) 651-8558</u>

### **Biorbyt LLC**

68 TW Alexander Drive Research Triangle Park Durham

NC 27713 United States

Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558





**Dilution range** 

ELISA: 1:10,000 - 1:50,000, IHC: 1:500 - 1:2,500, WB: 1:1,000 - 1:10,000

**Application notes** 

Anti-Sheep IgG F(c) Peroxidase Conjugated Antibody has been tested by ELISA and is assayed against 1.0 ug of Sheep IgG in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) as a substrate for 30 minutes at room temperature. A working dilution of 1:2,000 to 1:8,000 of the reconstitution concentration is suggested for this product.

**Antibody Type** 


Secondary Antibody

Storage

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

**Note** 

For research use only



ELISA Results of Rabbit Anti-Sheep IgG F(c) Antibody Peroxidase Conjugated tested against purified Sheep IgG F(c) HRP. Each well was coated in duplicate with 1.0  $\mu$ g of Sheep IgG F(c) (p/n orb346368). The working dilution is 1:35800. The starting dilution of antibody was 5  $\mu$ g/ml and the X-axis represents the Log10 of a 3-fold dilution. This titration is a 4-parameter curve fit where the IC50 is defined as the titer of the antibody. Assay performed using TMB substrate (p/n orb348651).

### **Biorbyt Ltd.**

7 Signet Court, Swann Road Cambridge CB5 8LA United Kingdom

Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u>
Phone: <u>+44 (0)1223 859353</u> | Fax: <u>+1 (415) 65</u>1-8558

68 TW Alexander Drive Research Triangle Park Durham NC 27713 United States

Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558