

## **Product Datasheet**

## Sheep IgG (H&L) Antibody Peroxidase Conjugated Pre-Adsorbed (orb347953)

Catalog Number orb347953

**Description** Sheep IgG (H&L) antibody (Peroxidase)

Species/Host Donkey

**Reactivity** Sheep

Conjugation HRP

**Tested Applications** ELISA, IHC, WB

Immunogen Anti-Sheep IgG (H&L) was produced by repeated immunization with Sheep IgG

whole molecule in donkey.

**Preservatives** 0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!

Form/Appearance Lyophilized

**Concentration** 1.0 mg/mL

**Storage** Store vial at 4° C prior to restoration. For extended storage aliquot contents and

freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to

immediate use.

**Note** For research use only

Application notes Anti-Sheep IgG (H&L) peroxidase conjugated antibody has been tested by ELISA

and is suitable for immunoblotting (western or dot blot), ELISA,

immunoperoxidase electron microscopy and immunohistochemistry as well as

other peroxidase-antibody based enzymatic assays requiring lot-to-lot

consistency.





**Isotype** IgG

**Clonality** Polyclonal

**Antibody Type** Secondary Antibody


**Purity** Anti-Sheep IgG (H&L) peroxidase conjugated antibody was prepared from

monospecific antiserum by immunoaffinity chromatography using Sheep IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Donkey Serum, Sheep IgG and Sheep Serum. No reaction was observed against Chicken, Guinea Pig, Hamster, Horse,

Human, Mouse, Rabbit and Rat Serum Proteins.

**Dilution Range** ELISA: 1:65,000, IHC: 1:500 - 1:2,500, WB: 1:1,000 - 1:10,000

**Expiration Date** 12 months from date of receipt.



ELISA results of purified Donkey Anti-Sheep IgG peroxidase conjugated MX8 Antibody tested against purified Sheep IgG. Each well was coated in duplicate with 1.0  $\mu$ g of Sheep IgG (green line). The starting dilution of antibody was 5  $\mu$ g/ml and the X-axis represents the Log10 of a 3-fold dilution. This titration is a 4-parameter curve fit where the IC50 is defined as the titer of the antibody.