

## **Product Datasheet**

## F(ab')2 Dog IgG (H&L) Antibody Biotin Conjugated (orb348103)

**Description** F(ab')2 Dog IgG (H&L) antibody (Biotin)

Species/Host Rabbit

**Reactivity** Canine

**Conjugation** Biotin

**Tested Applications** ELISA, IHC, WB

**Immunogen** Dog IgG whole molecule

**Preservatives** 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Form/Appearance Lyophilized

**Concentration** 1 mg/mL

**Storage** Store vial at 4° C prior to restoration. For extended storage aliquot contents and

freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to

immediate use.

**Note** For research use only

**Application notes** F(ab')2 Anti-Dog IgG Biotin Antibody has been tested by ELISA and is designed

for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.

**Isotype** IgG F(ab')2

**Clonality** Polyclonal






## **Antibody Type** Secondary Antibody

**Purity** This product was prepared from monospecific antiserum by immunoaffinity

chromatography using Dog IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Biotin, anti-Rabbit Serum, Dog IgG and Dog Serum. No reaction was observed against anti-Pepsin or anti-Rabbit IgG F(c).

**Dilution Range** ELISA: 1:20,000 - 1:100,000, IHC: 1:1,000-1:5,000, WB: 1:2,000-1:10,000

**Expiration Date** 12 months from date of receipt.



ELISA results of purified Rabbit F(ab')2 Anti-DOG IgG Antibody Biotin Conjugated tested against purified Dog IgG. Each well was coated in duplicate with 1.0  $\mu$ g of Dog IgG (Blue Line). The starting dilution of antibody was 5  $\mu$ g/ml and the X-axis represents the Log10 of a 3-fold dilution. This titration is a 4-parameter curve fit where the IC50 is defined as the titer of the antibody. Assay performed using Blocking buffer, Streptavidin-HRP conjugated (p/n orb348767), and TMB substrate (p/n orb348651).