

Product Datasheet

F(Ab')2 Fragment Of Hamster IgG (H&L) Antibody Biotin Conjugated Pre-Adsorbed (orb348407)

Description	F(Ab')2 Fragment Of Hamster IgG (H&L) antibody (Biotin)
Species/Host	Goat
Reactivity	Golden Hamster, Hamster
Conjugation	Biotin
Tested Applications	ELISA, IHC, WB
Immunogen	Golden Syrian and Armenian Hamster IgG whole molecules
Preservatives	0.01% (w/v) Sodium Azide
Form/Appearance	Lyophilized
Concentration	1 mg/mL
Storage	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Note	For research use only
Application notes	F(ab')2 Anti-Golden Syrian & Armenian Hamster IgG Biotin Antibody has been tested by ELISA and is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.
lsotype	IgG F(ab')2
Clonality	Polyclonal

Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: <u>+44 (0) 1223 859-353</u> | Fax: <u>+1 (415) 651-8558</u>

Biorbyt LLC.

68 TW Alexander Drive, Durham, NC, 27713, United States Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>




Biorbyt.com

Antibody TypeSecondary AntibodyPurityThis product was prepared from monospecific antiserum by immunoaffinity
chromatography using Golden Syrian and Armenian Hamster IgG coupled to
agarose beads followed by solid phase adsorption(s) to remove any unwanted
reactivities, pepsin digestion and chromatographic separation. Assay by
immunoelectrophoresis resulted in a single precipitin arc against anti-biotin,
anti-Goat Serum, Golden Syrian and Armenian Hamster IgG and Golden Syrian
and Armenian Hamster Serum. No reaction was observed against anti-Pepsin,
anti-Goat IgG F(c) and Mouse or Rat Serum Proteins.Dilution RangeELISA: 1:5000 - 1:25,000, IHC: 1:300-1:2000, WB: 1:2,000-1:10,000

Expiration Date

12 months from date of receipt.



ELISA results of purified F(ab')2 Fragment of GOAT Anti-GOLDEN SYRIAN & ARMENIAN HAMSTER IgG Biotin Conjugated Antibody tested against purified GOLDEN SYRIAN & ARMENIAN HAMSTER IgG. Each well was coated in duplicate with 1.0 μ g of antigen. The starting dilution of antibody was 5 μ g/ml and the X-axis represents the Log10 of a 3-fold dilution. This titration is a 4parameter curve fit where the IC50 is defined as the titer of the antibody. Assay performed using Blocking buffer, Streptavidin HRP conjugate 1:10000.

Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: +44 (0) 1223 859-353 | Fax: +1 (415) 651-8558

Biorbyt LLC.

68 TW Alexander Drive, Durham, NC, 27713, United States Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>