

## **Product Datasheet**

## Fab Mouse IgG (H&L) Antibody Texas Red Conjugated (orb348475)

**Description** Fab Mouse IgG (H&L) antibody (Texas Red)

Species/Host Goat

**Reactivity** Mouse

**Conjugation** Texas Red

**Tested Applications** DOT, FC, FLISA, IF

**Immunogen** Mouse IgG whole molecule

**Preservatives** 0.01% (w/v) Sodium Azide

Form/Appearance Lyophilized

**Concentration** 1.0 mg/mL

**Storage** Store vial at 4° C prior to restoration. For extended storage aliquot contents and

freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to

immediate use.

**Note** For research use only

**Application notes** Fab Anti-Mouse IgG (H&L) Texas Red Antibody has been tested by dot blot and is

suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. This product is also suitable for multiplex analysis, including

multicolor imaging, utilizing various commercial platforms.

**Isotype** IgG Fab





**Clonality** Polyclonal

Antibody Type Secondary Antibody


**Purity** This product was prepared from monospecific antiserum by immunoaffinity

chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, papain digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum. No reaction was observed against

anti-Papain or anti-Goat IgG F(c).

**Dilution Range** FLISA: 1:10,000 - 1:50,000, FC: 1:500 - 1:2,500, IF: 1:1,000 - 1:5,000

**Expiration Date** 12 months from date of receipt.



Dot Blot of Fab Anti-MOUSE IgG (GOAT) Texas Red™ Conjugated Antibody. Lane 1: 100 ng. Lane 2: 33.3 ng. Lane 3: 11.1 ng. Lane 4: 3.7 ng. Lane 5: 1.23 ng. Secondary Antibody: Goat anti-Mouse IgG Texas Red. Blocking Buffer: BlockOut orb348644 for 30 min at RT.