

## Product Datasheet

### Anti-F4/80 [Cl:A3-1 (recombinant version)] (orb411544)

<b>Description</b>	Rat monoclonal antibody to F4/80
<b>Species/Host</b>	Rat
<b>Reactivity</b>	Mouse
<b>Conjugation</b>	Unconjugated
<b>Tested Applications</b>	FC, IHC
<b>Immunogen</b>	Thioglycollate stimulated peritoneal macrophages of mouse origin.
<b>Target</b>	F4/80
<b>Preservatives</b>	PBS with 0.02% Proclin 300.
<b>Concentration</b>	1 mg/ml
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Note</b>	For research use only
<b>Isotype</b>	Rat IgG
<b>Clonality</b>	Monoclonal
<b>Clone Number</b>	Cl:A3-1 (recombinant version)
<b>Antibody Type</b>	Recombinant Antibody
<b>Purity</b>	Purified
<b>Uniprot ID</b>	<b>Q61549</b>

#### Biorbyt Ltd.


7 Signet Court, Swann's Road,  
Cambridge, CB5 8LA, United Kingdom  
Email: [info@biorbyt.com](mailto:info@biorbyt.com), [support@biorbyt.com](mailto:support@biorbyt.com)  
Phone: [+44 \(0\) 1223 859-353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)

#### Biorbyt LLC.

68 TW Alexander Drive,  
Durham, NC, 27713, United States  
Email: [info@biorbyt.com](mailto:info@biorbyt.com), [support@biorbyt.com](mailto:support@biorbyt.com)  
Phone: [+1 \(415\) 906-5211](tel:+1(415)9065211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)

**Expiration Date**

12 months from date of receipt.




Competitive flow-cytometry assay between anti-F4/80 CI:A3-1 variants and an existing commercial anti-F4/80 antibody. Mouse (*Mus musculus*) splenocytes were labelled ex vivo with a commercially available APC-labelled anti-F4/80 antibody and APE labelled anti-CD11b antibody and subject to flow-cytometry analysis (A), in which a small subpopulation of F4/80-CD11b positive cells may be observed. Subsets of commercial anti-F4/80 antibody-labelled splenocytes were also subsequently incubated with unlabelled versions of either the rat (*Rattus norvegicus*) IgG2b chimeric version (orb411543, B) or mouse IgG2A chimeric (C) version of CI:A3-1. Loss of the F4/80-CD11b positive subpopulation may be observed, demonstrating displacement of the commercial antibody and the specificity of CI:A3-1.

**Biorbyt Ltd.**

7 Signet Court, Swann's Road,  
Cambridge, CB5 8LA, United Kingdom  
Email: [info@biorbyt.com](mailto:info@biorbyt.com), [support@biorbyt.com](mailto:support@biorbyt.com)  
Phone: [+44 \(0\) 1223 859-353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)

**Biorbyt LLC.**

68 TW Alexander Drive,  
Durham, NC, 27713, United States  
Email: [info@biorbyt.com](mailto:info@biorbyt.com), [support@biorbyt.com](mailto:support@biorbyt.com)  
Phone: [+1 \(415\) 906-5211](tel:+1(415)9065211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)




In Figure 1 murine bone marrow-derived macrophages (BMDMs) were pre-blocked with rat anti-mouse CD16 & CD32 (clone FCR-4G8) and stained with non-recombinant anti-F4/80 [Cl:A3-1] conjugated to Alexa Fluor® 647 (AF647), all commercially available from competitors. In Figure 2 BMDMs were stained with recombinant anti-F4/80 [Cl:A3-1] or isotype control (anti-fluorescein [4-4-20 (enhanced)] IgG2b (orb256396). In Figure 3 BMDMs were stained with Fc Silent™ recombinant anti-F4/80 [Cl:A3-1] or isotype control (Fc Silent™ anti-fluorescein [4-4-20 (enhanced)] IgG2b. These were fluorescently labelled using the secondary antibody, goat IgG anti-rat IgG (H&L-chain) polyclonal antibody directly conjugated to Alexa Fluor® 647 (AF647) commercially available from a competitor. Whilst in Figure 2 the highest fluorescence signal is seen with the recombinant anti-F4/80 IgG2 (the isotype of the original hybridoma-derived antibody), the isotype control IgG2b (orb256396) shows considerable signal overlap, indicative of binding of the antibody to Fc-receptors. This illustrates the importance of isotype controls in such experiments when using conventional antibody formats particularly when Fc-blocking reagents are incompatible with the system used due to reactivity with the secondary antibody. The Fc silent™ format however overcomes this issue as seen in Figure 3, where the Fc silent™ recombinant anti-F4/80 yields a strong and distinct signal, whilst the isotype control shows no discernible difference to the background staining from the secondary antibody alone. Therefore, with Fc Silent™ reagents, no Fc-blocking products are required.

#### Biorbyt Ltd.

7 Signet Court, Swann's Road,  
Cambridge, CB5 8LA, United Kingdom  
Email: [info@biorbyt.com](mailto:info@biorbyt.com), [support@biorbyt.com](mailto:support@biorbyt.com)  
Phone: [+44 \(0\) 1223 859-353](tel:+44101223859353) | Fax: [+1 \(415\) 651-8558](tel:+14156518558)

#### Biorbyt LLC.

68 TW Alexander Drive,  
Durham, NC, 27713, United States  
Email: [info@biorbyt.com](mailto:info@biorbyt.com), [support@biorbyt.com](mailto:support@biorbyt.com)  
Phone: [+1 \(415\) 906-5211](tel:+14159065211) | Fax: [+1 \(415\) 651-8558](tel:+14156518558)



In Figure 1 murine bone marrow-derived macrophages (BMDMs) were pre-blocked with rat anti-mouse CD16 & CD32 (clone FCR-4G8) and stained with non-recombinant anti-F4/80 [Cl:A3-1] conjugated to Alexa Fluor® 647 (AF647), all commercially available from competitors. In Figure 2 BMDMs were stained with recombinant anti-F4/80 [Cl:A3-1] or isotype control (anti-fluorescein [4-4-20 (enhanced)] IgG2b (orb256396). In Figure 3 BMDMs were stained with Fc Silent™ recombinant anti-F4/80 [Cl:A3-1] or isotype control (Fc Silent™ anti-fluorescein [4-4-20 (enhanced)] IgG2b. These were fluorescently labelled using the secondary antibody, goat IgG anti-rat IgG (H&L-chain) polyclonal antibody directly conjugated to Alexa Fluor® 647 (AF647) commercially available from a competitor. Whilst in Figure 2 the highest fluorescence signal is seen with the recombinant anti-F4/80 IgG2 (the isotype of the original hybridoma-derived antibody), the isotype control IgG2b (orb256396) shows considerable signal overlap, indicative of binding of the antibody to Fc-receptors. This illustrates the importance of isotype controls in such experiments when using conventional antibody formats particularly when Fc-blocking reagents are incompatible with the system used due to reactivity with the secondary antibody. The Fc silent™ format however overcomes this issue as seen in Figure 3, where the Fc silent™ recombinant anti-F4/80 yields a strong and distinct signal, whilst the isotype control shows no discernible difference to the background staining from the secondary antibody alone. Therefore, with Fc Silent™ reagents, no Fc-blocking products are required.

#### Biorbyt Ltd.

7 Signet Court, Swann's Road,  
Cambridge, CB5 8LA, United Kingdom  
Email: [info@biorbyt.com](mailto:info@biorbyt.com), [support@biorbyt.com](mailto:support@biorbyt.com)  
Phone: [+44 \(0\) 1223 859-353](tel:+44101223859353) | Fax: [+1 \(415\) 651-8558](tel:+14156518558)

#### Biorbyt LLC.

68 TW Alexander Drive,  
Durham, NC, 27713, United States  
Email: [info@biorbyt.com](mailto:info@biorbyt.com), [support@biorbyt.com](mailto:support@biorbyt.com)  
Phone: [+1 \(415\) 906-5211](tel:+14159065211) | Fax: [+1 \(415\) 651-8558](tel:+14156518558)