



## Product Datasheet UBE2I UBC9 Antibody (orb443231)

Catalog Number orb443231

**Category** Antibodies

**Description** Anti-UBE2I UBC9 Antibody. Tested in Flow Cytometry, IF, IHC, IHC-F, ICC, WB

applications. This antibody reacts with Human, Mouse, Rat.

**Clonality** Polyclonal

Species/Host Rabbit

**Isotype** Rabbit IgG

**Conjugation** Unconjugated

**Reactivity** Human, Mouse, Rat

Form/Appearance Lyophilized

**Concentration** Adding 0.2 ml of distilled water will yield a concentration of 500 µg/ml.

**Purification** Immunogen affinity purified.

**Immunogen** A synthetic peptide corresponding to a sequence at the C-terminus of human

UBE2I UBC9, identical to the related mouse and rat sequences.

UniProt ID P63279

MW 18 kDa

**Tested applications** FC, ICC, IF, IHC, IHC-Fr, WB

**Application notes** Western blot, 0.1-0.5μg/ml Immunohistochemistry (Paraffin-embedded Section),

0.5-1µg/ml Immunohistochemistry (Frozen Section), 0.5-1µg/ml

 $Immunocytochemistry/Immunofluorescence, \ 2\mu g/ml \ Flow \ Cytometry \ (Fixed), \ 1-$ 

 $3\mu g/1x106$  cells. Add 0.2ml of distilled water will yield a concentration of

500ug/ml






**Cross Reactivity** No cross-reactivity with other proteins.

**Antibody Type** Primary Antibody


**Storage** Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -

20°C in small aliquots to prevent freeze-thaw cycles.

**Note** For research use only




Flow Cytometry analysis of A431 cells using anti-UBE2I/UBC9 antibody. Overlay histogram showing A431 cells (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-UBE2I/UBC9 Antibody (1  $\mu$ g/1x10^6 cells) for 30 min at 20°C. DyLight® 488 conjugated goat anti-rabbit IgG (5-10  $\mu$ g/1x10^6 cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1  $\mu$ g/1x10^6) used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.




IF analysis of UBE2I/UBC9 using anti-UBE2I/UBC9 antibody. UBE2I/UBC9 was detected in immunocytochemical section of U20S cell. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 2  $\mu$ g/mL rabbit anti-UBE2I/UBC9 Antibody overnight at 4°C. DyLight® 488 Conjugated Goat Anti-Rabbit IgG was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.








IHC analysis of UBE2I UBC9 using anti-UBE2I UBC9 antibody. UBE2I UBC9 was detected in paraffin-embedded section of human appendicitis tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1  $\mu$ g/ml rabbit anti-UBE2I UBC9 Antibody overnight at 4°C. Biotinylated goat antirabbit lgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) with DAB as the chromogen.



IHC analysis of UBE2I UBC9 using anti-UBE2I UBC9 antibody. UBE2I UBC9 was detected in paraffin-embedded section of human gastric cancer tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1  $\mu$ g/ml rabbit anti-UBE2I UBC9 Antibody overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) with DAB as the chromogen.




IHC analysis of UBE2I UBC9 using anti-UBE2I UBC9 antibody. UBE2I UBC9 was detected in paraffin-embedded section of human glioma tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1  $\mu$ g/ml rabbit anti-UBE2I UBC9 Antibody overnight at 4°C. Biotinylated goat anti-rabbit lgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) with DAB as the chromogen.


Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>








IHC analysis of UBE2I UBC9 using anti-UBE2I UBC9 antibody. UBE2I UBC9 was detected in paraffin-embedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1  $\mu$ g/ml rabbit anti-UBE2I UBC9 Antibody overnight at 4°C. Biotinylated goat antirabbit lgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) with DAB as the chromogen.




IHC analysis of UBE2I UBC9 using anti-UBE2I UBC9 antibody. UBE2I UBC9 was detected in paraffin-embedded section of human placenta tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1  $\mu$ g/ml rabbit anti-UBE2I UBC9 Antibody overnight at 4°C. Biotinylated goat anti-rabbit lgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) with DAB as the chromogen.




IHC analysis of UBE2I UBC9 using anti-UBE2I UBC9 antibody. UBE2I UBC9 was detected in paraffin-embedded section of mouse brain tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1  $\mu$ g/ml rabbit anti-UBE2I UBC9 Antibody overnight at 4°C. Biotinylated goat antirabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) with DAB as the chromogen.








IHC analysis of UBE2I UBC9 using anti-UBE2I UBC9 antibody. UBE2I UBC9 was detected in paraffin-embedded section of rat brain tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1  $\mu$ g/ml rabbit anti-UBE2I UBC9 Antibody overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) with DAB as the chromogen.



IHC analysis of UBE2I UBC9 using anti-UBE2I UBC9 antibody. UBE2I UBC9 was detected in paraffin-embedded section of rat lung tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1  $\mu$ g/ml rabbit anti-UBE2I UBC9 Antibody overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) with DAB as the chromogen.




IHC analysis of UBE2I/UBC9 using anti-UBE2I/UBC9 antibody. UBE2I/UBC9 was detected in paraffin-embedded section of human rectal cancer tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1  $\mu$ g/ml rabbit anti-UBE2I/UBC9 Antibody overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) with DAB as the chromogen.


Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>








IHC analysis of UBE2I/UBC9 using anti-UBE2I/UBC9 antibody. UBE2I/UBC9 was detected in paraffin-embedded section of human thyroid cancer tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1  $\mu$ g/ml rabbit anti-UBE2I/UBC9 Antibody overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) with DAB as the chromogen.



IHC analysis of UBE2I/UBC9 using anti-UBE2I/UBC9 antibody. UBE2I/UBC9 was detected in paraffin-embedded section of human tonsil tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1  $\mu$ g/ml rabbit anti-UBE2I/UBC9 Antibody overnight at 4°C. Biotinylated goat antirabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) with DAB as the chromogen.





IHC analysis of UBE2I/UBC9 using anti-UBE2I/UBC9 antibody. UBE2I/UBC9 was detected in paraffin-embedded section of mouse intestine tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1  $\mu$ g/ml rabbit anti-UBE2I/UBC9 Antibody overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) with DAB as the chromogen.


Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558












IHC analysis of UBE2I/UBC9 using anti-UBE2I/UBC9 antibody. UBE2I/UBC9 was detected in paraffin-embedded section of rat intestine tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1  $\mu$ g/ml rabbit anti-UBE2I/UBC9 Antibody overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) with DAB as the chromogen.

Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>







Western blot analysis of UBE2I UBC9 using anti-UBE2I UBC9 antibody. Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human placenta tissue lysates, Lane 2: human K562 whole cell lysates, Lane 3: human HepG2 whole cell lysates, Lane 4: rat brain tissue lysates, Lane 5: rat kidney tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-UBE2I UBC9 antigen affinity purified polyclonal antibody at 0.5 μg/mL overnight at 4°C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for UBE2I UBC9 at approximately 18 kDa. The expected band size for UBE2I UBC9 is at 18 kDa.