

Product Datasheet

COX IV COX4I1 Antibody (monoclonal, 4G11) (orb548371)

Catalog Number orb548371

Category Antibodies

Description Anti-COX IV COX4I1 Antibody (monoclonal, 4G11). Tested in Flow Cytometry,

IHC, WB applications. This antibody reacts with Human, Mouse, Rat.

Clonality Monoclonal

Species/Host Mouse

Isotype Mouse IgG2b

Conjugation Unconjugated

Reactivity Human, Mouse, Rat

Form/Appearance Lyophilized

Concentration Adding 0.2 ml of distilled water will yield a concentration of 500 μg/ml.

Purification Immunogen affinity purified.

Immunogen E. coli-derived human COX IV recombinant protein (Position: Q59-K169).

UniProt ID P13073

MW 17 kDa

Tested applications FC, IHC, WB

Application notes Western blot, 0.1-0.5μg/ml Immunohistochemistry (Paraffin-embedded Section),

0.5-1µg/ml Flow Cytometry (Fixed), 1-3µg/1x106 cells. Add 0.2ml of distilled

water will yield a concentration of 500µg/ml





Cross Reactivity No cross-reactivity with other proteins.

Antibody Type Primary Antibody


Clone Number 4G11

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -


20°C in small aliquots to prevent freeze-thaw cycles.

Note For research use only

Expiration Date 12 months from date of receipt.




Flow Cytometry analysis of U937 cells using anti-COX IV antibody. Overlay histogram showing U937 cells (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-COX IV Antibody (1 $\mu g/1x10^6$ cells) for 30 min at 20°C. DyLight® 488 conjugated goat anti-mouse IgG (5-10 $\mu g/1x10^6$ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was mouse IgG (1 $\mu g/1x10^6$) used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.




IHC analysis of COX IV using anti-COX IV antibody. COX IV was detected in paraffin-embedded section of human colon cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 μ g/ml mouse anti-COX IV Antibody overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) with DAB as the chromogen.







IHC analysis of COX IV using anti-COX IV antibody. COX IV was detected in paraffin-embedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 μ g/ml mouse anti-COX IV Antibody overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) with DAB as the chromogen.




IHC analysis of COX IV using anti-COX IV antibody. COX IV was detected in paraffin-embedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 μ g/ml mouse anti-COX IV Antibody overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) with DAB as the chromogen.

$$\label{eq:phone:mail:model} \begin{split} &\text{Email: } \underline{info@biorbyt.com}, \ \underline{support@biorbyt.com} \\ &\text{Phone: } \underline{+1 \ (415) \ 906-5211} \ \big| \ \text{Fax: } \underline{+1 \ (415) \ 651-8558} \end{split}$$







Western blot analysis of COX IV using anti-COX IV antibody. Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50 ug of sample under reducing conditions. Lane 1: human HEPG2 whole cell lysates, Lane 2: human A549 whole cell lysates, Lane 3: human HEK293 whole cell lysates, Lane 4: human T47D whole cell lysates, Lane 5: human CACO-2 whole cell lysates, Lane 6: human K562 whole cell lysates, Lane 7: human Hela whole cell lysates, Lane 8: rat brain tissue lysates, Lane 9: mouse brain tissue lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-COX IV antigen affinity purified monoclonal antibody at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for COX IV at approximately 17KD. The expected band size for COX IV is at 17KD.

Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>