

Product Datasheet

MUC4 Antibody / Mucin-4 (orb607148)

Description The major constituents of mucus, the viscous secretion that covers epithelial

surfaces such as those in the trachea, colon, and cervix, are highly glycosylated

proteins called mucins. These glycoproteins play important roles in the

and differentiation. This gene encodes an integral membrane glycoprotein found on the cell surface, although secreted isoforms may exist. At least two dozen transcript variants of this gene have been found, although for many of them the

protection of the epithelial cells and have been implicated in epithelial renewal

full-length transcript has not been determined or they are found only in tumor

tissues. [RefSeq]

Species/Host Mouse

Reactivity Human

Conjugation Unconjugated

Tested Applications IHC-P

Immunogen Amino acids 1730-1864 from the human protein were used as the immunogen

for this Mucin 4 antibody.

Preservatives 0.2 mg/ml in 1X PBS with 0.1 mg/ml rAlbumin (US sourced) and 0.05% sodium

azide

Storage Store the Mucin 4 antibody at 2-8°C (with azide) or aliquot and store at -20°C or

colder (without azide).

Note For research use only

Application notes Optimal dilution of the Mucin 4 antibody should be determined by the

researcher.1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb

solution onto the tissue section and incubate at RT for 30 min.

Formula 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide

Isotype Mouse IgG1, kappa

Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>





Clonality Monoclonal

Clone Number MUCN4-1

Antibody Type Primary Antibody

Purity Protein G affinity chromatography


Uniprot ID Q99102

Hazard Information This Mucin 4 antibody is available for research use only.

Dilution Range Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT,Prediluted IHC only

format: incubate for 30 min at RT (1)

Expiration Date 12 months from date of receipt.



IHC testing of FFPE human colon carcinoma with Mucin 4 antibody (clone MUCN4-1). HIER: requires steaming of sections in pH9 10mM Tris with 1mM EDTA for 10-20 min and allow to cool before testing.

Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558