



## Product Datasheet MCM6 Antibody (orb639801)

Catalog Number orb639801

**Category** Antibodies

**Description** The mini-chromosome maintenance (MCM) family of proteins, including MCM2,

MCM3, MCM6 (Cdc21), MCM5 (Cdc46), MCM6 (Mis5) and MCM7 (Cdc47), are regulators of DNA replication that act to ensure replication occurs only once in the cell cycle. Expression of MCM proteins increases during cell growth, peaking at G1/S phase. The MCM proteins each contain an ATP-binding motif, which is predicted to mediate ATP-dependent opening of double-stranded DNA. MCM proteins are regulated by E2F transcription factors, which induce MCM

expression, and by protein kinases, which interact with MCM proteins to maintain

the post-replicative state of the cell. MCM2/MCM6 complexes function as

substrates for Cdc2/cyclin B in vitro.

**Clonality** Monoclonal

**Species/Host** Mouse

**Isotype** Mouse IgG2b, kappa

**Conjugation** Unconjugated

**Reactivity** Human

**Immunogen** A recombinant human partial protein (amino acids 228-368) was used as the

immunogen for this MCM6 antibody.

UniProt ID Q14566

**Tested applications** IHC-P

**Dilution range** Immunohistochemistry (FFPE): 1-2ug/ml

**Application notes** Optimal dilution of the antibody should be determined by the researcher.

Antibody Type Primary Antibody






Clone Number MCM6/2999

**Formula** 0.2 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide


**Storage** Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -

20°C in small aliquots to prevent freeze-thaw cycles.

**Note** For research use only




IHC staining of FFPE human tonsil with MCM6 antibody. HIER: boil tissue sections in pH9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human colon with MCM6 antibody. HIER: boil tissue sections in pH9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.









IHC staining of FFPE human colon carcinoma with MCM6 antibody. HIER: boil tissue sections in pH9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

Human Protein Microarray Specificity Validation



Analysis of HuProt (TM) microarray containing more than 19000 full-length human proteins using MCM6 antibody (clone MCM6/2999). These results demonstrate the foremost specificity of the MCM6/2999 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt (TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt (TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



SDS-PAGE analysis of purified, BSA-free MCM6 antibody as confirmation of integrity and purity.