



## **Product Datasheet**

## Cd79a Antibody (orb654317)

Catalog Number orb654317

**Category** Antibodies

**Description** Anti-Cd79a Antibody. Tested in ELISA, Flow Cytometry, IF, IHC, WB applications.

This antibody reacts with Mouse, Rat.

**Clonality** Polyclonal

Species/Host Rabbit

**Isotype** Rabbit IgG

**Conjugation** Unconjugated

**Reactivity** Mouse, Rat

Form/Appearance Lyophilized

**Concentration** Adding 0.2 ml of distilled water will yield a concentration of 500 µg/ml.

**Purification** Immunogen affinity purified.

**Immunogen** E.coli-derived mouse Cd79a recombinant protein (Position: L29-P220).

UniProt ID P11911

**MW** 25-30 kDa

**Tested applications** ELISA, FC, IF, IHC, WB

**Application notes** Western blot, 0.25-0.5μg/ml, Mouse, Rat Immunohistochemistry (Paraffin-

embedded Section), 0.5-1 $\mu$ g/ml, Mouse, Rat Immunofluorescence, 2 $\mu$ g/ml, Mouse, Rat Flow Cytometry (Fixed), 1-3 $\mu$ g/1x106 cells, Mouse ELISA, 0.1-0.5 $\mu$ g/ml, -. Add 0.2ml of distilled water will yield a concentration of 500 $\mu$ g/ml

**Cross Reactivity** No cross-reactivity with other proteins.

**Biorbyt Ltd.** 

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom Email: <a href="mailto:info@biorbyt.com">info@biorbyt.com</a> Phone: +44 (0) 1223 859-353 | Fax: +1 (415) 651-8558 **Biorbyt LLC.** 

68 TW Alexander Drive,

Durham, NC, 27713, United States
Email: <a href="mailto:info@biorbyt.com">info@biorbyt.com</a>, <a href="mailto:support@biorbyt.com">support@biorbyt.com</a>

Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558






**Antibody Type** Primary Antibody

**Storage** Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -


20°C in small aliquots to prevent freeze-thaw cycles.

**Note** For research use only

**Expiration Date** 12 months from date of receipt.




Flow Cytometry analysis of mouse spleen tissues using anti-Cd79a antibody. Overlay histogram showing mouse spleen tissues (Blue line). The tissues were fixed with 4% paraformaldehyde and blocked with 10% normal goat serum. And then incubated with rabbit anti-Cd79a Antibody (1  $\mu g/1x10^6$  cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (5-10  $\mu g/1x10^6$  cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1  $\mu g/1x10^6$ ) used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.




IF analysis of Cd79a using anti-Cd79a antibody. Cd79a was detected in paraffin-embedded section of mouse intestine (lymph node) tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2  $\mu$ g/mL rabbit anti-Cd79a Antibody overnight at 4°C. DyLight®488 Conjugated Goat Anti-Rabbit IgG was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.








IF analysis of Cd79a using anti-Cd79a antibody. Cd79a was detected in paraffin-embedded section of rat intestine tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2  $\mu$ g/mL rabbit anti-Cd79a Antibody overnight at 4°C. DyLight®488 Conjugated Goat Anti-Rabbit IgG was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.




IHC analysis of Cd79a using anti-Cd79a antibody. Cd79a was detected in paraffin-embedded section of mouse thymus tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1  $\mu$ g/ml rabbit anti-Cd79a Antibody overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) with DAB as the chromogen.



IHC analysis of Cd79a using anti-Cd79a antibody. Cd79a was detected in paraffin-embedded section of rat thymus tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1  $\mu$ g/ml rabbit anti-Cd79a Antibody overnight at 4°C. Biotinylated goat anti-rabbit lgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) with DAB as the chromogen.







Western blot analysis of Cd79a using anti-Cd79a antibody. Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50 ug of sample under reducing conditions. Lane 1: rat spleen tissue lysates, Lane 2: mouse spleen tissue lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Cd79a antigen affinity purified polyclonal antibody at 0.5 μg/mL overnight at 4°C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for Cd79a at approximately 25-30 kDa. The expected band size for Cd79a is at 25 kDa.

Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558