



Product Datasheet UBE2L6 Antibody (orb668554)

Catalog Number orb668554

Category Antibodies

Description The UBE2L6 Antibody is suitable for IF, WB. It is a Polyclonal, Unconjugated

antibody which raised against KLH-conjugated synthetic peptide of human UBE2L6. The exact sequence is proprietary. Purification: The antibody was

purified by immunogen affinity chromatography.

Clonality Polyclonal

Species/Host Rabbit

Conjugation Unconjugated

Reactivity Human, Mouse, Rat

Form/Appearance Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30%

glycerol, and 0.01% sodium azide.

Purification The antibody was purified by immunogen affinity chromatography.

Immunogen KLH-conjugated synthetic peptide of human UBE2L6. The exact sequence is

proprietary.

UniProt ID Q9QZU9, Q4V8J2, O14933

Tested applications IF, WB

Dilution range WB: 1-500-2000

Antibody Type Primary Antibody

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -

20°C in small aliquots to prevent freeze-thaw cycles.

Note For research use only

Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom Email: info@biorbyt.com Phone: +44 (0) 1223 859-353 | Fax: +1 (415) 651-8558 **Biorbyt LLC.**

68 TW Alexander Drive, Durham, NC, 27713, United States

Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558






Entrez


56791, 295704, 9246

Expiration Date

12 months from date of receipt.



Western blot analysis of UBE2L6 expression in Jurkat (A), mouse lung (B), mouse spleen (C), rat kidney (D), rat liver (E) whole cell lysates. (Predicted band size: 10; 17 kD; Observed band size: 18 kD)



Immunofluorescent analysis of UBE2L6 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AF594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>