

Product Datasheet

Chicken anti-Goat IgG (H&L), TRITC conjugated (orb700060)

Catalog Number	orb700060
Description	Gallus polyclonal antibody against Gallus anti-Goat IgG (H&L) conjugated to TRITC
Species/Host	Gallus
Reactivity	Goat
Conjugation	TRITC
Form/Appearance	Liquid
Storage	Store at 2-8°C. Shelf life of this product is one year from date of receipt.
Note	For research use only
Application notes	This antibody reacts with the heavy chains on goat IgG and with the light chains on all goat immunoglobulins based on immunoelectrophoresis.Minimum cross- reactivity is observed to non-immunoglobulin goat serum proteins based on immunoelectrophoresis.Antibody is supplied in 10 mM sodium phosphate, 150 mM sodium chloride, pH 7.2, 1% (w/v) BSA, protease/IgG free and 0.05 % (w/v) sodium azide as preservative.BSA and milk have to be replaced by other blocking reagents, like doneky serum or commercial formulations which are free from bovine IgG.
Application notes	on all goat immunoglobulins based on immunoelectrophoresis.Minimum cross- reactivity is observed to non-immunoglobulin goat serum proteins based on immunoelectrophoresis.Antibody is supplied in 10 mM sodium phosphate, 150 mM sodium chloride, pH 7.2, 1% (w/v) BSA, protease/IgG free and 0.05 % (w/v) sodium azide as preservative.BSA and milk have to be replaced by other blocking reagents, like doneky serum or commercial formulations which are free
	on all goat immunoglobulins based on immunoelectrophoresis.Minimum cross- reactivity is observed to non-immunoglobulin goat serum proteins based on immunoelectrophoresis.Antibody is supplied in 10 mM sodium phosphate, 150 mM sodium chloride, pH 7.2, 1% (w/v) BSA, protease/IgG free and 0.05 % (w/v) sodium azide as preservative.BSA and milk have to be replaced by other blocking reagents, like doneky serum or commercial formulations which are free from bovine IgG.
Clonality	on all goat immunoglobulins based on immunoelectrophoresis.Minimum cross- reactivity is observed to non-immunoglobulin goat serum proteins based on immunoelectrophoresis.Antibody is supplied in 10 mM sodium phosphate, 150 mM sodium chloride, pH 7.2, 1% (w/v) BSA, protease/IgG free and 0.05 % (w/v) sodium azide as preservative.BSA and milk have to be replaced by other blocking reagents, like doneky serum or commercial formulations which are free from bovine IgG. Polyclonal

Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: <u>+44 (0) 1223 859-353</u> | Fax: <u>+1 (415) 651-8558</u>

Biorbyt LLC.

68 TW Alexander Drive, Durham, NC, 27713, United States Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>