

Product Datasheet

Donkey anti-Goat IgG (H&L), ALP conjugated (orb700094)

Description	Donkey polyclonal antibody against Donkey anti-Goat IgG (H&L) conjugated to AP
Species/Host	Donkey
Conjugation	ALP
Form/Appearance	Liquid
Storage	Non-diluted antibody is stable for 4 years at 2-8°C. For storage at -20°C dilute antibody solution with an equal volume of glycerol to obtain final glycerol concentration of 50 % to prevent loss of enzymatic activity. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard, Be sure to mix well but without foaming.
Note	For research use only
Application notes	This antibody reacts with the heavy chains on goat IgG and with the light chains on all goat immunoglobulins.Minimum cross-reactivity is observed to non- immunoglobulin goat serum proteins based on immunoelectrophoresis.Antibody is supplied in 30 mM triethanolamine, pH 7.2, 5 mM magnesium chloride, 0.1 mM zinc chloride, 1 % (w/v) BSA, Protease/IgG free , 0.05 % (w/v) sodium azide as preservative.BSA and milk have to be replaced by other blocking reagents, like doneky serum or commercial formulations which are free from bovine IgG.
Clonality	Polyclonal
Purity	Immunogen affinity purified IgG.
Dilution Range	1 : 50-1 : 5000 (ICC), 1 : 20-1 : 2000 (IHC), 1 : 500-1 : 2000 (WB)
Expiration Date	12 months from date of receipt.

Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: <u>+44 (0) 1223 859-353</u> | Fax: <u>+1 (415) 651-8558</u>

Biorbyt LLC.

68 TW Alexander Drive, Durham, NC, 27713, United States Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>