

Product Datasheet Anti-CD31 Antibody (orb705400)

Description Mouse monoclonal antibody to CD31

Species/Host Mouse

Reactivity Human

Conjugation Unconjugated

Tested Applications IH, WB

Immunogen KLH-conjugated synthetic peptide encompassing a sequence within human

CD31. The exact sequence is proprietary.

Target PECAM1

Preservatives Mouse IgG1. Liquid in PBS containing 50% glycerol, 0.2% rAlbumin and 0.01%

sodium azide.

Form/Appearance Mouse IgG1. Liquid in PBS containing 50% glycerol, 0.2% rAlbumin and 0.01%

sodium azide.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -

20°C in small aliquots to prevent freeze-thaw cycles.

Note For research use only

Clonality Monoclonal

Antibody Type Primary Antibody

Source Mouse

Uniprot ID P16284

Entrez 5175

Dilution Range WB: 1:500-1:1000, IHC-P: 1:100-1:300

Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom Email: info@biorbyt.com Phone: +44 (0) 1223 859-353 | Fax: +1 (415) 651-8558 **Biorbyt LLC.**

68 TW Alexander Drive,

Durham, NC, 27713, United States


Email: $\underline{info@biorbyt.com}$, $\underline{support@biorbyt.com}$ Phone: $\underline{+1 (415) 906-5211}$ | Fax: $\underline{+1 (415) 651-8558}$






Expiration Date


12 months from date of receipt.



Western blot analysis of CD31 expression in MDAMB231 (A), Hela (B) whole cell lysates. (Predicted band size: 82 kD; Observed band size: 130 kD)



Immunohistochemical analysis of CD31 staining in human hemangioendothelioma formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.




Immunohistochemical analysis of CD31 staining in human placenta formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.


Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>







Immunohistochemical analysis of CD31 staining in human tonsil formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunohistochemical analysis of CD31 staining in human liver formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.