

Product DatasheetPSD95/DLG4 Antibody (orb745945)

Catalog Number orb745945

Category Antibodies

Description Anti-PSD95/DLG4 Antibody. Tested in ELISA, Flow Cytometry, WB applications.

This antibody reacts with Human, Mouse, Rat.

Clonality Polyclonal

Species/Host Rabbit

Isotype Rabbit IgG

Conjugation Unconjugated

Reactivity Human, Mouse, Rat

Form/Appearance Lyophilized

Concentration Adding 0.2 ml of distilled water will yield a concentration of 500 μg/ml.

Purification Immunogen affinity purified.

Immunogen E.coli-derived human PSD95/DLG4 recombinant protein (Position: D18-L711).

UniProt ID P78352

MW 99 kDa

Tested applications ELISA, FC, WB

Application notes Western blot, 0.25-0.5μg/ml, Mouse, Rat Flow Cytometry (Fixed), 1-3μg/1x106

cells, Human ELISA, 0.1-0.5µg/ml, -. Add 0.2ml of distilled water will yield a

concentration of 500ug/ml

Biorbyt Ltd.

7 Signet Court, Swann Road Cambridge CB5 8LA United Kingdom

Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: +44 (0)1223 859353 | Fax: +1 (415) 651-8558

Biorbyt LLC

68 TW Alexander Drive Research Triangle Park

Durham NC 27713 United States

Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558






Cross Reactivity No cross-reactivity with other proteins.

Antibody Type Primary Antibody

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -

20°C in small aliquots to prevent freeze-thaw cycles.

Note For research use only



Flow Cytometry analysis of U937 cells using anti-PSD95/DLG4 antibody. Overlay histogram showing U937 cells (Blue line). The cells were fixed with 4% paraformaldehyde and blocked with 10% normal goat serum. And then incubated with rabbit anti-PSD95/DLG4 Antibody (1 μ g/1x10^6 cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (5-10 μ g/1x10^6 cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 μ g/1x10^6) used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.

Biorbyt Ltd.

7 Signet Court, Swann Road Cambridge CB5 8LA United Kingdom

Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u>
Phone: +44 (0)1223 859353 | Fax: +1 (415) 651-8558


Biorbyt LLC

68 TW Alexander Drive Research Triangle Park Durham NC 27713 United States

Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558







Western blot analysis of PSD95/DLG4 using anti-PSD95/DLG4 antibody. Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: rat brain tissue lysates, Lane 2: mouse brain tissue lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-PSD95/DLG4 antigen affinity purified polyclonal antibody at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for PSD95/DLG4 at approximately 99 KD. The expected band size for PSD95/DLG4 is at 99 KD.

Biorbyt Ltd.

7 Signet Court, Swann Road Cambridge CB5 8LA United Kingdom

Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: +44 (0)1223 859353 | Fax: +1 (415) 651-8558

Biorbyt LLC

68 TW Alexander Drive Research Triangle Park Durham NC 27713 United States

Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558