

Product Datasheet

Goat Anti-Rabbit IgG H&L, Cy3 conjugated (orb868589)

| | |
|-----------------------------|---|
| Catalog Number | orb868589 |
| Category | Antibodies |
| Description | The Goat Anti-Rabbit IgG H&L, Cy3 conjugated is a Cy3 conjugated polyclonal antibody suitable for FC, ICC, IF. This product reacts with Rabbit samples. |
| Clonality | Polyclonal |
| Species/Host | Goat |
| Isotype | IgG |
| Conjugation | Cy3 |
| Reactivity | Rabbit |
| Form/Appearance | Liquid |
| Concentration | 2.0 mg/ml |
| Buffer/Preservatives | 10 mM TBS (pH=7.4) with 1% rAlbumin, 0.03% Proclin300 and 50% glycerol. |
| Purification | Affinity purified by Protein G, nonspecific adsorbed |
| Immunogen | Native rabbit IgG |
| Tested applications | FC, ICC, IF |
| Antibody Type | Secondary Antibody |
| Storage | Store at -20°C for one year. Avoid repeated freeze/thaw cycles. |

Biorbyt Ltd.

7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom

Email: info@biorbyt.com, support@biorbyt.com

Phone: [+44 \(0\)1223 859353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)

Biorbyt LLC


68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713-2847
United States

Email: info@biorbyt.com, support@biorbyt.com


Phone: [+1 \(415\) 906-5211](tel:+1(415)906-5211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)

Note For research use only


Expiration Date 12 months from date of receipt.



Paraformaldehyde-fixed, paraffin embedded (human lung carcinoma), Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min, Blocking buffer (normal goat serum) at 37°C for 30 min, Incubation with (c-Maf) Polyclonal Antibody, Unconjugated at 1:200 overnight at 4°C, followed by a conjugated Goat Anti-Rabbit IgG antibody for 90 minutes, and DAPI for nuclei staining.



Paraformaldehyde-fixed, paraffin embedded (mouse brain), Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min, Blocking buffer (normal goat serum) at 37°C for 30 min, Incubation with (c-Maf) Polyclonal Antibody, Unconjugated at 1:200 overnight at 4°C, followed by a conjugated Goat Anti-Rabbit IgG antibody for 90 minutes, and DAPI for nuclei staining.



Paraformaldehyde-fixed, paraffin embedded (rat brain), Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min, Blocking buffer (normal goat serum) at 37°C for 30 min, Incubation with (c-Maf) Polyclonal Antibody, Unconjugated at 1:200 overnight at 4°C, followed by a conjugated Goat Anti-Rabbit IgG antibody for 90 minutes, and DAPI for nuclei staining.

Biorbyt Ltd.

7 Signet Court, Swann Road
Cambridge
CB5 8LA
United Kingdom

Email: info@biorbyt.com, support@biorbyt.com
Phone: [+44 \(0\)1223 859353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)

Biorbyt LLC

68 TW Alexander Drive
Research Triangle Park
Durham
NC 27713-2847
United States

Email: info@biorbyt.com, support@biorbyt.com
Phone: [+1 \(415\) 906-5211](tel:+1(415)906-5211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)651-8558)